

AD-A189 707

THE PARTIALLY OBSERVED STOCHASTIC MINIMUM PRINCIPLE(U)  
ALBERTA UNIV EDMONTON DEPT OF STATISTICS AND APPLIED  
PROBABILITY J BARAS ET AL. 11 NOV 87 AFOSR-TR-87-1939

1/1


UNCLASSIFIED

AFOSR-86-0332

F/G 12/3

ML





FILE COPY

②

AD-A189 787

## REPORT DOCUMENTATION PAGE

2a. SECURITY CLASSIFICATION AUTHORITY			1b. RESTRICTIVE MARKINGS			
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release: distribution unlimited.			
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 87 - 1939</b>			
6a. NAME OF PERFORMING ORGANIZATION University of Alberta		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM			
6c. ADDRESS (City, State, and ZIP Code) Edmonton, Alberta, Canada T6G 2G1			7b. ADDRESS (City, State, and ZIP Code) <b>AFOSR/NM</b> Bldg 410 Bolling AFB DC 20332-6448			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (if applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-86-0332			
8c. ADDRESS (City, State, and ZIP Code) Bolling AFB DC 20332-6448			10. SOURCE OF FUNDING NUMBERS			
			PROGRAM ELEMENT NO 61102F	PROJECT NO 2304	TASK NO A1	WORK UNIT ACCESSION NO
11. TITLE (Include Security Classification) The Partially Observed Stochastic Minimum Principle						
12. PERSONAL AUTHOR(S) John Baras, Robert J. Elliott & Michael Kohlmann						
13a. TYPE OF REPORT Reprint		13b. TIME COVERED FROM 9/30/86 TO 9/30/87		14. DATE OF REPORT (Year, Month, Day) Nov. 11, 1987		15. PAGE COUNT
16. SUPPLEMENTARY NOTATION						
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD	GROUP	SUB-GROUP				
19. ABSTRACT (Continue on reverse if necessary and identify by block number)						
<p>The focus of this research is the filtering jump processes. To investigate the filtering of manifold-valued processes, their approximation by random walks and Markov chains was studied. The object was to approximate a signal process by a finite-state jump process for which a finite-dimensional filter is available. Four papers were published during the past year, including "The existence of smooth densities for the prediction, filtering and smoothing problems" and "The partially observed stochastic minimum principle".</p>						
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION <b>E</b>			
22a. NAME OF RESPONSIBLE INDIVIDUAL Maj James Crowley			22b. TELEPHONE (Include Area Code) (202) 767-5025		22c. OFFICE SYMBOL NM	

UNCLASSIFIED

DTIC  
ELECTE  
JAN 05 1988  
ca

**AFOSR-TR. 87-1939**

**The Partially Observed Stochastic  
Minimum Principle**

JOHN BARAS  
UNIVERSITY OF MARYLAND, USA  
ROBERT J. ELLIOTT  
UNIVERSITY OF ALBERTA, CANADA  
MICHAEL KOHLMANN  
UNIVERSITÄT KONSTANZ, F.R. GERMANY

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



12 May 1987

87 12 23 247

**ACKNOWLEDGEMENTS.** The second and third authors wish to thank the Natural Sciences and Engineering Research Council of Canada for its support under grant A-7964. The second author was partially supported by the Air Force Office of Scientific Research, United States Air Force, under grant AFOSR-86-0332 and European Office of Aerospace Research and Development, London, England.

# **The Partially Observed Stochastic Minimum Principle**

JOHN BARAS  
SYSTEMS RESEARCH CENTER, UNIVERSITY OF MARYLAND  
COLLEGE PARK, MD 20742 USA

ROBERT J. ELLIOTT  
DEPARTMENT OF STATISTICS AND APPLIED PROBABILITY, UNIVERSITY OF ALBERTA  
EDMONTON, AB T6G 2G1 CANADA

MICHAEL KOHLMANN  
FAKULTAT FÜR WIRTSCHAFTSWISSENSCHAFTEN UND STATISTIK  
UNIVERSITÄT KONSTANZ, POSTFACH 5560, D-7750 F.R. GERMANY

## **1. INTRODUCTION.**

Various proofs have been given of the minimum principle satisfied by an optimal control in a partially observed stochastic control problem. See, for example, the papers by Bensoussan [1], Elliott [5], Haussmann [7], and the recent paper [9] by Haussmann in which the adjoint process is identified. The simple case of a partially observed Markov chain is discussed in the University of Maryland lecture notes [6] of the second author.

We show in this article how a minimum principle for a partially observed diffusion can be obtained by differentiating the statement that a control  $u^*$  is optimal. The results of Bismut [2], [3] and Kunita [10], on stochastic flows enable us to compute in an easy and explicit way the change in the cost due to a 'strong variation' of an optimal control. The only technical difficulty is the justification of the differentiation. As we wished to exhibit the simplification obtained by using the ideas of stochastic flows the result is not proved under the weakest possible hypotheses. Finally, in Section 6, we show how Bensoussan's minimum principle follows from our result if the drift coefficient is differentiable in the control variable.

## 2. DYNAMICS.

Suppose the state of the system is described by a stochastic differential equation

$$\begin{aligned} d\xi_t &= f(t, \xi_t, u)dt + g(t, \xi_t)dw_t, \\ \xi_t &\in R^d, \quad \xi_0 = x_0, \quad 0 \leq t \leq T. \end{aligned} \quad (2.1)$$

The control parameter  $u$  will take values in a compact subset  $U$  of some Euclidean space  $R^k$ .

We shall make the following assumptions:

$A_1$ :  $x_0$  is given; if  $x_0$  is a random variable and  $P_0$  its distribution the situation when  $\int |x|^q P_0(dx) < \infty$  for some  $q > n + 1$  can be treated, as in [9], by including an extra integration with respect to  $P_0$ .

$A_2$ :  $f : [0, T] \times R^d \times U \rightarrow R^d$  is Borel measurable, continuous in  $u$  for each  $(t, x)$ , continuously differentiable in  $x$  and for some constant  $K$

$$(1 + |x|)^{-1} |f(t, x, u)| + |f_x(t, x, u)| \leq K_1.$$

$A_3$ :  $g : [0, T] \times R^d \rightarrow R^d \otimes R^n$  is a matrix valued function, Borel measurable, continuously differentiable in  $x$ , and for some constant  $K_2$

$$|g(t, x)| + |g_x(t, x)| \leq K_2.$$

The observation process is given by

$$\begin{aligned} dy_t &= h(\xi_t)dt + dv_t \\ y_t &\in R^m, \quad y_0 = 0, \quad 0 \leq t \leq T. \end{aligned} \quad (2.2)$$

In the above equations  $w = (w^1, \dots, w^n)$  and  $v = (v^1, \dots, v^d)$  are independent Brownian motions. We also assume

$A_4$ :  $h : R^d \rightarrow R^m$  is Borel measurable, continuously differentiable in  $x$ , and for some constant  $K_3$

$$|h(t, x)| + |h_x(t, x)| \leq K_3.$$

REMARKS 2.1. These hypotheses can be weakened. For example, in  $A_4$ ,  $h$  can be allowed linear growth in  $x$ . Because  $g$  is bounded a delicate argument then implies the exponential  $Z$  of (2.3) is in some  $L^p$  space,  $1 < p < \infty$ . (See, for example, Theorem 2.2 of [8]). However, when  $h$  is bounded  $Z$  is in all the  $L^p$  spaces, (see Lemma 2.3). Also, if we require  $f$  to have linear growth in  $u$  then the set of control values  $U$  can be unbounded as in [9]. Our objective, however, is not the greatest generality but to demonstrate the simplicity of the techniques of stochastic flows.

Let  $\hat{P}$  denote Wiener measure on the  $C([0, T], R^n)$  and  $\mu$  denote Wiener measure on  $C([0, T], R^m)$ . Consider the space  $\Omega = C([0, T], R^n) \times C([0, T], R^m)$  with coordinate functions  $(x_t, y_t)$  and define Wiener measure  $P$  on  $\Omega$  by

$$P(dx, dy) = \hat{P}(dx)\mu(dy).$$

DEFINITION 2.2. Write  $Y = \{Y_t\}$  for the right continuous complete filtration on  $C([0, T], R^m)$  generated by  $Y_t^0 = \sigma\{y_s : s \leq t\}$ . The set of admissible control functions  $\underline{U}$  will be the  $Y$ -predictable functions on  $[0, T] \times C([0, T], R^m)$  with values in  $U$ .

For  $u \in \underline{U}$  and  $x \in R^d$  write  $\xi_{s,t}^u(x)$  for the strong solution of (2.1) corresponding to control  $u$ , and with  $\xi_{s,s}^u(x) = x$ . Write

$$Z_{s,t}^u(x) = \exp \left( \int_s^t h(\xi_{s,r}^u(x))' dy_r - \frac{1}{2} \int_s^t h(\xi_{s,r}^u(x))^2 dt \right) \quad (2.3)$$

and define a new probability measure  $P^u$  on  $\Omega$  by  $\frac{dP^u}{dP} = Z_{0,T}^u(x_0)$ . Then under  $P^u$   $(\xi_{0,t}^u(x_0), y_t)$  is a solution of (2.1) and (2.2), that is  $\xi_{0,t}^u(x_0)$  remains a strong solution of (2.1) and there is an independent Brownian motion  $v$  such that  $y_t$  satisfies (2.2). A version of  $Z$  defined for every trajectory  $y$  of the observation process is obtained by integrating by parts the stochastic integral in (2.3).

LEMMA 2.3. Under hypothesis  $A_4$ , for  $t \leq T$ ,

$$E[(Z_{0,t}^u(x_0))^p] < \infty \quad \text{for all } u \in \underline{U} \text{ and all } p, \quad 1 \leq p < \infty.$$



PROOF.

$$Z_{0,t}^u(x_0) = 1 + \int_0^t Z_{0,r}^u(x_0) h(\xi_{0,r}^u(x_0))' dy_r.$$

Therefore, for any  $p$  there is a constant  $C_p$  such that

$$E[(Z_{0,t}^u(x_0))^p] \leq C_p \left[ 1 + E\left(\int_0^t (Z_{0,r}^u(x_0))^2 h(\xi_{0,r}^u(x_0))^2 dr\right)^{p/2} \right].$$

The result follows by Gronwall's inequality.

COST 2.4. We shall suppose the cost is purely terminal and given by some bounded, differentiable function

$$c(\xi_{0,T}^u(x_0))$$

which has bounded derivatives. Then the expected cost if control  $u \in \underline{U}$  is used is

$$J(u) = E_u[c(\xi_{0,T}^u(x_0))].$$

In terms of  $P$ , under which  $y_t$  is always a Brownian motion, this is

$$J(e) = E\left[Z_{0,T}^u(x_0) c(\xi_{0,T}^u(x_0))\right]. \quad (2.4)$$

### 3. STOCHASTIC FLOWS.

For  $u \in \underline{U}$  write

$$\xi_{s,t}^u(x) = x + \int_s^t f(r, \xi_{s,r}^u(x), u_r) dr + \int_s^t g(r, \xi_{s,r}^u(x)) dw_r \quad (3.1)$$

for the solution of (2.1) over the time interval  $[s, t]$  with initial condition  $\xi_{s,s}^u(x) = x$ . In the sequel we wish to discuss the behaviour of (3.1) for each trajectory  $y$  of the observation process. We have already noted there is a version of  $Z$  defined for every  $y$ . The results of Bismut [2] and Kunita [10] extend easily and show the map

$$\xi_{s,t}^u : R^d \rightarrow R^d$$

is, almost surely, for each  $y \in C([0, T], R^m)$  a diffeomorphism. Bismut [2] initially gives proofs when the coefficients  $f$  and  $g$  are bounded, but points out that a stopping time argument extends the results to when, for example, the coefficients have linear growth.

Write  $\|\xi^u(x_0)\|_t = \sup_{0 \leq s \leq t} |\xi_{0,s}^u(x_0)|$ . Then, as in Lemma 2.1 of [8], for any  $p$ ,  $1 \leq p < \infty$  using Gronwall's and Jensen's inequalities

$$\|\xi^u(x_0)\|_T^p \leq C \left( 1 + |x_0|^p + \left| \int_0^T g(r, \xi_{0,r}^u(x_0)) dw_r \right|^p \right)$$

almost surely, for some constant  $C$ .

Therefore, using Burkholder's inequality and hypothesis  $A_3$ ,  $\|\xi^u(x_0)\|_T$  is in  $L^p$  for all  $p$ ,  $1 \leq p < \infty$ .

Suppose  $u^* \in U$  is an optimal control so  $J(u^*) \leq J(u)$  for any other  $u \in U$ . Write  $\xi_{s,t}^*(\cdot)$  for  $\xi_{s,t}^{u^*}(\cdot)$ . The Jacobian  $\frac{\partial \xi_{s,t}^*}{\partial x}(x)$  is the matrix solution  $C_t$  of the equation for  $s \leq t$ ,

$$dC_t = f_x(t, \xi_{s,t}^*(x), u^*) C_t dt + \sum_{i=1}^n g_x^{(i)}(t, \xi_{s,t}^*(x)) C_t dw_t^i \quad (3.2)$$

with  $C_s = I$ .

Here  $I$  is the  $n \times n$  identity matrix and  $g^{(i)}$  is the  $i^{\text{th}}$  column of  $g$ . From hypotheses  $A_2$  and  $A_3$ ,  $f_x$  and  $g_x$  are bounded. Writing  $\|C\|_T = \sup_{0 \leq s \leq T} |C_s|$  an application of Gronwall's

Jensen's and Burkholder's inequalities again implies  $\|C\|_T$  is in  $L^p$  for all  $p$ ,  $1 \leq p < \infty$ .

Consider the related matrix valued stochastic differential equation

$$\begin{aligned} D_t = I - \int_s^t D_r f_x(r, \xi_{s,r}^*(x), u_r^*)' dr \\ - \sum_{i=1}^n \int_s^t D_r g_x^{(i)}(r, \xi_{s,r}^*(x))' dw_r^i \\ + \sum_{i=1}^n \int_s^t D_r (g_x^{(i)}(r, \xi_{s,r}^*(x)))'^2 dr. \end{aligned} \quad (3.3)$$

Then it can be checked that  $D_t C_t = I$  for  $t \geq s$ , so that  $D_t$  is the inverse of the Jacobian, that is  $D_t = \left( \frac{\partial \xi_{s,t}^*(x)}{\partial x} \right)^{-1}$ . Again, because  $f_x$  and  $g_x$  are bounded we have that  $\|D\|_t$  is in every  $L^p$ ,  $1 \leq p < \infty$ .

For a  $d$ -dimensional semimartingale  $z_t$  Bismut [2] shows one can consider the flow  $\xi_{s,t}^*(z_t)$  and gives the semimartingale representation of this process. In fact if  $z_t = z_s + A_t + \sum_{i=1}^n \int_s^t H_i dw_r^i$  is the  $d$ -dimensional semimartingale, Bismut's formula states that

$$\begin{aligned} \xi_{s,t}^*(z_t) = z_s + \int_s^t \left( f(r, \xi_{s,r}^*(z_r), u_r^*) \right. \\ + \sum_{i=1}^n g_x^{(i)}(r, \xi_{s,r}^*(z_r), u_r^*) \frac{\partial \xi_{s,r}^*(z_r)}{\partial x} H_i + \frac{1}{2} \sum_{i=1}^n \frac{\partial^2 \xi_{s,r}^*(z_r)}{\partial x^2} (H_i, H_i) \Big) dr \\ + \int_s^t \frac{\partial \xi_{s,r}^*(z_r)}{\partial x} dA_r + \sum_{i=1}^n \int_s^t \left( g^{(i)}(r, \xi_{s,r}^*(z_r)) + \frac{\partial \xi_{s,r}^*(z_r)}{\partial x} H_i \right) dw_r^i. \end{aligned} \quad (3.4)$$

**DEFINITION 3.1.** We shall consider perturbations of the optimal control  $u^*$  of the following kind: For  $s \in [0, T)$ ,  $h > 0$  such that  $0 \leq s < s + h \leq T$ , for any other admissible control  $u \in \underline{U}$  and  $A \in Y_s$  define a strong variation of  $u^*$  by

$$u(t, w) = \begin{cases} u^*(t, w) & \text{if } (t, w) \notin [s, s+h] \times A \\ \tilde{u}(t, w) & \text{if } (t, w) \in [s, s+h] \times A. \end{cases}$$

Applying (3.4) as in Theorem 5.1 of [4] we have the following result.

**THEOREM 3.2.** For the perturbation  $u$  of the optimal control  $u^*$  consider the process

$$z_t = x + \int_s^t \left( \frac{\partial \xi_{s,r}^*(z_r)}{\partial x} \right)^{-1} \left( f(r, \xi_{s,r}^*(z_r), u_r) - f(r, \xi_{s,r}^*(z_r), u_r^*) \right) dr. \quad (3.5)$$

Then the process  $\xi_{s,t}^*(z_t)$  is indistinguishable from  $\xi_{s,t}^u(x)$ .

PROOF. Note the equation defining  $z_t$  involves only an integral in time; there is no martingale term, so to apply (3.4) we have  $H_i = 0$  for all  $i$ . Therefore, from (3.4)

$$\begin{aligned}\xi_{s,t}^*(z_t) &= x + \int_s^t f(r, \xi_{s,r}^*(z_r), u_r^*) dr \\ &\quad + \int_s^t \left( \frac{\partial \xi_{s,r}^*(z_r)}{\partial x} \right) \left( \frac{\partial \xi_{s,r}^*(z_r)}{\partial x} \right)^{-1} (f(r, \xi_{s,r}^*(z_r), u_r) - f(r, \xi_{s,r}^*(z_r), u_r^*)) dr \\ &\quad + \int_s^t g(r, \xi_{s,r}^*(z_r)) dw_r.\end{aligned}$$

However, the solution of (3.2) is unique so

$$\xi_{s,t}^*(z_t) = \xi_{s,t}^u(x).$$

REMARKS 3.3. Note that the perturbation  $u(t)$  equals  $u^*(t)$  if  $t > s+h$  so  $z_t = z_{s+h}$  if  $t > s+h$  and

$$\xi_{s,r}^*(z_r) = \xi_{s,t}^*(z_{s+h}) = \xi_{s+h,t}^*(\xi_{s,s+h}^u(x)).$$

#### 4. AUGMENTED FLOWS.

Consider the augmented flow which includes as an extra coordinate the stochastic exponential  $Z_{s,t}^*$  with a 'variable' initial condition  $z \in R$  for  $Z_{s,s}^*(\cdot)$ . That is, consider the  $(d+1)$  dimensional system given by:

$$\begin{aligned}\xi_{s,t}^*(x) &= x + \int_s^t f(r, \xi_{s,r}^*(x), u_r^*) dr + \int_s^t g(r, \xi_{s,r}^*(x)) dw_r \\ Z_{s,t}^*(x, z) &= z + \int_s^t Z_{s,r}^*(x, z) h(\xi_{s,r}^*(x))' dy_r.\end{aligned}$$

Therefore,

$$\begin{aligned}Z_{s,t}^*(x, z) &= z Z_{s,r}^*(x) \\ &= z \exp \left( \int_s^t h(\xi_{s,r}^*(x))' dy_r - \frac{1}{2} \int_s^t h(\xi_{s,r}^*(x))^2 dr \right)\end{aligned}$$

and we see there is a version of the enlarged system defined for each trajectory  $y$  by integrating by parts the stochastic integral. The augmented map  $(x, z) \rightarrow (\xi_{s,t}^*(x), Z_{s,t}^*(x, z))$  is then almost surely a diffeomorphism of  $R^{d+1}$ . Note that  $\frac{\partial \xi_{s,t}^*(x)}{\partial z} = 0$ ,  $\frac{\partial f}{\partial z} = 0$  and  $\frac{\partial g}{\partial z} = 0$ . The Jacobian of this augmented map is, therefore, represented by the matrix

$$\tilde{C}_t = \begin{pmatrix} \frac{\partial \xi_{s,t}^*(x)}{\partial z} & 0 \\ \frac{\partial Z_{s,t}^*(x, z)}{\partial z} & \frac{\partial Z_{s,t}^*(x, z)}{\partial z} \end{pmatrix},$$

and for  $1 \leq i \leq d$  from equation (3.3)

$$\begin{aligned}\frac{\partial Z_{s,t}^*(x, z)}{\partial x_i} &= \sum_{j=1}^m \int_s^t \left( Z_{s,r}^*(x, z) \frac{\partial h^j(\xi_{s,r}^*(x))}{\partial \xi_k} \cdot \frac{\partial \xi_{k,s,r}^*(x)}{\partial x_i} \right. \\ &\quad \left. + h^j(\xi_{s,r}^*(x)) \frac{\partial Z_{s,r}^*(x, z)}{\partial x_i} \right) dy_r^j.\end{aligned}\tag{4.1}$$

(Here the double index  $k$  is summed from 1 to  $n$ ).

We shall be interested in the solution of this differential system (4.1) only in the situation when  $z = 1$  so we shall write  $Z_{s,t}^*(x)$  for  $Z_{s,t}^*(x, 1)$ . The following result is motivated by formally differentiating the exponential formula for  $Z_{s,t}^*(x)$ .

LEMMA 4.1.

$$\frac{\partial Z_{s,t}^*(x)}{\partial x} = Z_{s,t}^*(x) \left( \int_s^t h_x(\xi_{s,r}^*(x)) \cdot \frac{\partial \xi_{s,r}^*(x)}{\partial x} \cdot dv_r \right)$$

where  $v = (v^1, \dots, v^n)$  is the Brownian motion in the observation process.

PROOF. From (4.1) we see  $\frac{\partial Z_{s,t}^*(x)}{\partial x}$  is the solution of the stochastic differential equation

$$\frac{\partial Z_{s,t}^*(x)}{\partial x} = \int_s^t \left( \frac{\partial Z_{s,r}^*(x)}{\partial x} h'(\xi_{s,r}^*(x)) + Z_{s,r}^*(x) h_x(\xi_{s,r}^*(x)) \frac{\partial \xi_{s,r}^*(x)}{\partial x} \right) dy_r. \quad (4.2)$$

Write

$$L_{s,t}(x) = Z_{s,t}^*(x) \left( \int_s^t h_x \cdot \frac{\partial \xi_{s,r}^*(x)}{\partial x} \cdot dv_r \right)$$

where

$$dy_r = h(\xi_{s,t}^*(x)) dt + dv_t.$$

Because

$$Z_{s,t}^*(x) = 1 + \int_s^t Z_{s,r}^*(x) h'(\xi_{s,r}^*(x)) dy_r$$

the product rule gives

$$\begin{aligned} L_{s,t}(x) &= \int_s^t Z_{s,r}^*(x) h_x \cdot \frac{\partial \xi_{s,r}^*(x)}{\partial x} dv_r \\ &\quad + \int_s^t \left( \int_s^r h_x \cdot \frac{\partial \xi_{s,\sigma}^*(x)}{\partial x} \cdot dv_\sigma \right) Z_{s,r}^*(x) h'(\xi_{s,r}^*(x)) dy_r \\ &\quad + \int_s^t Z_{s,r}^*(x) h'(\xi_{s,r}^*(x)) \cdot h_x \cdot \frac{\partial \xi_{s,r}^*(x)}{\partial x} dr \\ &= \int_s^t L_{s,r}(x) h'(\xi_{s,r}^*(x)) dy_r + \int_s^t Z_{s,r}^*(x) h_x \cdot \frac{\partial \xi_{s,r}^*(x)}{\partial x} \cdot dy_r. \end{aligned}$$

Therefore,  $L_{s,t}(x)$  is also a solution of (4.2) so by uniqueness

$$L_{s,t}(x) = \frac{\partial Z_{s,t}^*(x)}{\partial x}.$$

REMARKS 4.2. As noted at the beginning of this section we can consider the augmented flow

$$(x, z) \rightarrow (\xi_{s,t}^*(x), Z_{s,t}^*(x, z)) \quad \text{for } x \in R^d, z \in R,$$

and we are only interested in the situation when  $z = 1$ , so we write  $Z_{s,t}^*(x)$ .

LEMMA 4.3.  $Z_{s,t}^*(z_t) = Z_{s,t}^u(x)$  where  $z_t$  is the semimartingale defined in (3.6).

PROOF.  $Z_{s,t}^u(x)$  is the process uniquely defined by

$$Z_{s,t}^u(x) = 1 + \int_s^t Z_{s,r}^u(x) h'(\xi_{s,r}^u(x)) dy_r. \quad (4.2)$$

Consider an augmented  $(d+1)$  dimensional version of (3.6) defining a semimartingale  $\bar{z}_t = (z_t, 1)$ , so the additional component is always identically 1. Then applying (3.5) to the new component of the augmented process we have

$$\begin{aligned} Z_{s,r}^*(z_r) &= 1 + \int_s^r Z_{s,r}^*(z_r) h'(\xi_{s,r}^*(z_r)) dy_r \\ &= 1 + \int_s^r Z_{s,r}^*(z_r) h'(\xi_{s,r}^u(x)) dy_r \end{aligned}$$

by Theorem 3.2. However, (4.2) has a unique solution so  $Z_{s,t}^*(z_t) = Z_{s,t}^u(x)$ .

REMARKS 4.4. Note that for  $t > s + h$

$$Z_{s,t}^*(z_t) = Z_{s,t}^*(z_{s+h}).$$

## 5. THE MINIMUM PRINCIPLE.

Control  $u$  will be the perturbation of the optimal control  $u^*$  as in Definition 3.1. We shall write  $x = \xi_{0,s}^*(x_0)$ . Then the minimum cost is

$$\begin{aligned} J(u^*) &= E[Z_{0,T}^*(x_0)c(\xi_{0,T}^*(x_0))] \\ &= E[Z_{0,s}^*(x_0)Z_{s,T}^*(x)c(\xi_{s,T}^*(x))]. \end{aligned}$$

The cost corresponding to the perturbed control  $u$  is

$$\begin{aligned} J(u) &= E[Z_{0,s}^*(x_0)Z_{s,T}^u(x)c(\xi_{s,T}^u(x))] \\ &= E[Z_{0,s}^*(x_0)Z_{s,T}^*(z_{s+h})c(\xi_{s,T}^*(z_{s+h}))] \end{aligned}$$

by Theorem 3.2 and Lemma 4.3. Now  $Z_{s,T}^*(\cdot)$  and  $c(\xi_{s,T}^*(\cdot))$  are differentiable with continuous and uniformly integrable derivatives. Therefore

$$\begin{aligned} J(u) - J(u^*) &= E[Z_{0,s}^*(x_0)(Z_{s,T}^*(z_{s+h})c(\xi_{s,T}^*(z_{s+h})) - Z_{s,T}^*(x)c(\xi_{s,T}^*(x)))] \\ &= E\left[\int_s^{s+h} \Gamma(s, z_r)(f(r, \xi_{s,r}^*(z_r), u_r) - f(r, \xi_{s,r}^*(x), u_r^*))dr\right] \end{aligned}$$

where

$$\begin{aligned} \Gamma(s, z_r) &= Z_{0,s}^*(x_0)Z_{s,T}^*(z_r)\left\{c_\xi(\xi_{s,T}^*(z_r))\frac{\partial \xi_{s,T}^*(z_r)}{\partial x} + \right. \\ &\quad \left. c(\xi_{s,T}^*(z_r))\left(\int_s^T h_\xi(\xi_{s,\sigma}^*(z_r))\frac{\partial \xi_{s,\sigma}^*(z_r)}{\partial x}dv_\sigma\right)\right\}\left(\frac{\partial \xi_{s,r}^*(z_r)}{\partial x}\right)^{-1}. \end{aligned}$$

Note that this expression gives an explicit formula for the change in the cost resulting from a variation in the optimal control. The only remaining problem is to justify differentiating the right hand side.

From Lemma 2.3,  $Z$  is in every  $L^p$  space,  $1 \leq p < \infty$  and from the remarks at the beginning of Section 3,  $C_T = \frac{\partial \xi_{s,T}^*}{\partial x}$  and  $D_T = \left(\frac{\partial \xi_{s,T}^*}{\partial x}\right)^{-1}$  are in every  $L^p$  space,  $1 \leq p < \infty$ . Consequently,  $\Gamma$  is in every  $L^p$  space,  $1 \leq p < \infty$ .



Therefore

$$\begin{aligned}
J(u) - J(u^*) &= \int_s^{s+h} E \left[ (\Gamma(s, z_r) - \Gamma(s, x)) (f(r, \xi_{s,r}^*(z_r), u_r) - f(r, \xi_{s,r}^*(z_r), u_r^*)) \right] dr \\
&\quad + \int_s^{s+h} E \left[ (\Gamma(s, x) - \Gamma(r, x)) (f(r, \xi_{s,r}^*(z_r), u_r) - f(r, \xi_{s,r}^*(z_r), u_r^*)) \right] dr \\
&\quad + \int_s^{s+h} E \left[ \Gamma(r, x) (f(r, \xi_{s,r}^*(z_r), u_r) - f(r, \xi_{s,r}^*(z_r), u_r^*) \right. \\
&\quad \quad \left. - f(r, \xi_{s,r}^*(x), u_r) + f(r, \xi_{s,r}^*(x), u_r^*)) \right] dr \\
&\quad + \int_s^{s+h} E \left[ \Gamma(r, x) (f(r, \xi_{0,r}^*(x_0), u_r) - f(r, \xi_{0,r}^*(x_0), u_r^*)) \right] dr \\
&= I_1(h) + I_2(h) + I_3(h) + I_4(h), \quad \text{say.}
\end{aligned}$$

Now,

$$\begin{aligned}
|I_1(h)| &\leq K_1 \int_s^{s+h} E \left[ |\Gamma(s, z_r) - \Gamma(s, x)| (1 + \|\xi^u(x_0)\|_{s+h}) \right] dr \\
&\leq K_1 h \sup_{s \leq r \leq s+h} E \left[ |\Gamma(s, z_r) - \Gamma(s, x)| (1 + \|\xi^u(x_0)\|_{s+h}) \right] \\
|I_2(h)| &\leq K_2 \int_s^{s+h} E \left[ |\Gamma(s, x) - \Gamma(r, x)| (1 + \|\xi^u(x_0)\|_{s+h}) \right] dr \\
&\leq K_2 h \sup_{s \leq r \leq s+h} E \left[ |\Gamma(s, z_r) - \Gamma(r, x)| (1 + \|\xi^u(x_0)\|_{s+h}) \right] \\
|I_3(h)| &\leq K_3 \int_s^{s+h} E \left[ |\Gamma(r, x)| \|x - z_r\| \right] dr \\
&\leq K_3 h \sup_{s \leq r \leq s+h} E \left[ |\Gamma(r, x)| \|x - z_r\|_{s+h} \right].
\end{aligned}$$

The differences  $|\Gamma(s, z_r) - \Gamma(s, x)|$ ,  $|\Gamma(s, x) - \Gamma(r, x)|$  and  $\|x - z_r\|_{s+h}$  are all uniformly bounded in some  $L^p$ ,  $p \geq 1$ , and

$$\lim_{r \rightarrow s} |\Gamma(s, z_r) - \Gamma(s, x)| = 0 \quad \text{a.s.}$$

$$\lim_{r \rightarrow s} |\Gamma(s, x) - \Gamma(r, x)| = 0 \quad \text{a.s.}$$

$$\lim_{h \rightarrow 0} \|x - z_r\|_{s+h} = 0.$$

Therefore,

$$\begin{aligned}\lim_{r \rightarrow s} \|\Gamma(s, z_r) - \Gamma(s, x)\|_p &= 0 \\ \lim_{r \rightarrow s} \|\Gamma(s, x) - \Gamma(r, x)\|_p &= 0 \\ \text{and } \lim_{h \rightarrow 0} \|(\|x - z\|_{s+h})\|_p &= 0 \text{ for some } p.\end{aligned}$$

Consequently,  $\lim_{h \rightarrow 0} h^{-1} I_k(h) = 0$ , for  $k = 1, 2, 3$ .

The only remaining problem concerns the differentiability of

$$I_4(h) = \int_s^{s+h} E \left[ \Gamma(r, x) (f(r, \xi_{0,r}^*(x_0), u_r) - f(r, \xi_{0,r}^*(x_0), u_r^*)) \right] dr.$$

The integrand is almost surely in  $L^1([0, T])$  so  $\lim_{h \rightarrow 0} h^{-1} I_4(h)$  exists for almost every  $s \in [0, T]$ . However, the set of times  $\{s\}$  where the limit may not exist might depend on the control  $u$ . Consequently we must restrict the perturbations  $u$  of the optimal control  $u^*$  to perturbations from a countable dense set of controls. In fact:

- 1) Because the trajectories are, almost surely, continuous,  $Y_\rho$  is countably generated by sets  $\{A_{i\rho}\}$ ,  $i = 1, 2, \dots$  for any rational number  $\rho \in [0, T]$ . Consequently  $Y_t$  is countably generated by the sets  $\{A_{i\rho}\}$ ,  $\rho \leq t$ .
- 2) Let  $G_t$  denote the set of measurable functions from  $(\Omega, Y_t)$  to  $U \subset R^k$ . (If  $u \in \underline{U}$  then  $u(t, w) \in G_t$ .) Using the  $L^1$ -norm, as in [5], there is a countable dense subset  $H_\rho = \{u_{j\rho}\}$  of  $G_\rho$ , for rational  $\rho \in [0, T]$ . If  $H_t = \bigcup_{\rho \leq t} H_\rho$  then  $H_t$  is a countable dense subset of  $G_t$ . If  $u_{j\rho} \in H_\rho$  then, as a function constant in time,  $u_{j\rho}$  can be considered as an admissible control over the time interval  $[t, T]$  for  $t \geq \rho$ .
- 3) The countable family of perturbations is obtained by considering sets  $A_{i\rho} \in Y_t$ , functions  $u_{j\rho} \in H_t$ , where  $\rho \leq t$ , and defining as in 3.1

$$u_{j\rho}^*(s, w) = \begin{cases} u^*(s, w) & \text{if } (s, w) \notin [t, T] \times A_{i\rho} \\ u_{j\rho}(s, w) & \text{if } (s, w) \in [t, T] \times A_{i\rho}. \end{cases}$$

Then for each  $i, j, \rho$

$$\lim_{h \rightarrow 0} h^{-1} \int_s^{s+h} E \left[ \Gamma(r, x) (f(r, \xi_{0,r}^*(x_0), u_{j\rho}^*) - f(r, \xi_{0,r}^*(x_0), u^*)) \right] dr \quad (5.1)$$

exists and equals

$$E \left[ \Gamma(s, x) (f(s, \xi_{0,s}^*(x_0), u_{j\rho}) - f(s, \xi_{0,s}^*(x_0), u^*)) I_{A_{j\rho}} \right]$$

for almost all  $s \in [0, T]$ .

Therefore, considering this perturbation we have

$$\begin{aligned} \lim_{h \rightarrow 0} h^{-1} (J(u_{j\rho}^*) - J(u^*)) &= E \left[ \Gamma(s, x) (f(s, \xi_{0,s}^*(x_0), u_{j\rho}) - f(s, \xi_{0,s}^*(x_0), u^*)) I_{A_{j\rho}} \right] \\ &\geq 0 \quad \text{for almost all } s \in [0, T]. \end{aligned}$$

Consequently there is a set  $S \subset [0, T]$  of zero Lebesgue measure such that, if  $s \notin S$ , the limit in (5.1) exists for all  $i, j, \rho$ , and gives

$$E \left[ \Gamma(s, x) (f(s, \xi_{0,s}^*(x_0), u_{j\rho}) - f(s, \xi_{0,s}^*(x_0), u^*)) I_{A_{j\rho}} \right] \geq 0.$$

Using the monotone class theorem, and approximating an arbitrary admissible control  $u \in \underline{U}$  we can deduce that if  $s \notin S$

$$E \left[ \Gamma(s, x) (f(s, \xi_{0,s}^*(x_0), u) - f(s, \xi_{0,s}^*(x_0), u^*)) I_A \right] \geq 0 \quad \text{for any } A \in Y_s. \quad (5.2)$$

Write

$$p_s(x) = E^* \left[ c_\xi(\xi_{0,T}^*(x_0)) \frac{\partial \xi_{s,T}^*(x)}{\partial x} + c(\xi_{0,T}^*(x_0)) \left( \int_s^T h_\xi(\xi_{0,\sigma}^*(x_0)) \frac{\partial \xi_{s,\sigma}^*(x)}{\partial x} dv_\sigma \right) \mid Y_s \vee \{x\} \right]$$

where, as before,  $x = \xi_{0,s}^*(x_0)$  and  $E^*$  denotes expectation under  $P^* = P^{u^*}$ . Then  $p_s(x)$  is the co-state variable and we have in (5.2) proved the following 'conditional' minimum principle:

**THEOREM 5.1.** *If  $u^* \in \underline{U}$  is an optimal control there is a set  $S \subset [0, T]$  of zero Lebesgue measure such that if  $s \notin S$*

$$E^* [p_s(x) f(s, x, u^*) \mid Y_s] \leq E^* [p_s(x) f(s, x, u) \mid Y_s] \quad \text{a.s.}$$

*That is, the optimal control  $u^*$  almost surely minimizes the conditional Hamiltonian and the adjoint variable is  $p_s(x)$ .*

## 6. CONCLUSION.

Using the theory of stochastic flows the effect of a perturbation of an optimal control is explicitly calculated. The only difficulty was to justify its differentiation. The adjoint process is explicitly identified as  $p_s(x)$ .

**THEOREM 6.1.** *If  $f$  is differentiable in the control variable  $u$ , and if the random variable  $x = \xi_{0,s}^*(x_0)$  has a conditional density  $q_s(x)$  under the measure  $P^*$ , then the inequality of Theorem 5.1 implies*

$$\sum_{j=1}^k (u_j(s) - u_j^*(s)) \int_{R^d} \Gamma(s, x) \frac{\partial f}{\partial u_j}(s, x, u^*) q_s(x) dx \leq 0.$$

This is the result of Bensoussan's paper [1].

The method of this paper can be applied to completely observable systems by initially considering 'stochastic open loop' controls, systems with stochastic constraints and deterministic systems. The adjoint process can be explicitly identified. 'Almost minimum' principles for 'almost optimal' controls can be obtained. Some of these will be discussed in later work.

# References

- [1] A. Bensoussan, *Maximum principle and dynamic programming approaches of the optimal control of partially observed diffusions*. Stochastics, 9(1983), 169–222.
- [2] J.M. Bismut, *A generalized formula of Ito and some other properties of stochastic flows*. Zeits. fur Wahrs. 55(1981), 331–350.
- [3] J.M. Bismut, *Mecanique Aléatoire*. Lecture Notes in Mathematics 866, Springer-Verlag, 1981.
- [4] J.M. Bismut, *Mecanique Aléatoire*. In Ecole d'Eté de Probabilités de Saint-Flour X. Lecture Notes in Mathematics 929, pp. 1–100. Springer-Verlag, 1982.
- [5] R.J. Elliott, *The optimal control of a stochastic system*. SIAM Jour. Control and Opt. 15(1977), 756–778.
- [6] R.J. Elliott, *Filtering and control for point process observations*. Systems Science Center, University of Maryland. Springer Lecture Notes, to appear.
- [7] U.G. Haussmann, *General necessary conditions for optimal control of stochastic systems*. Math. Programming Studies 6(1976), 30–48.
- [8] U.G. Haussmann, *A Stochastic Minimum Principle for Optimal Control of Diffusions*. Pitman Research Notes in Mathematics 151, Longman U.K., 1986.
- [9] U.G. Haussmann, *The maximum principle for optimal control of diffusions with partial information*. SIAM J. Control and Optimization 25(1987), 341–361.
- [10] H. Kunita, *On the decomposition of solutions of stochastic differential equations*. Lecture Notes in Mathematics 851(1980), 213–255.

END

DATE

FILMED

APRIL

1988

DTIC